

**PCT**WORLD INTELLECTUAL  
PROPERTY ORGANIZATION  
Inter

## INTERNATIONAL APPLICATION PUBLISHED

WO 9603151A2

(51) International Patent Classification <sup>6</sup> : <b>A61K 48/00 // C12N 9/48</b>		<b>A2</b>	(43) International Publication Date: 8 February 1996 (08.02.96)
(21) International Application Number: PCT/GB95/01783 (22) International Filing Date: 27 July 1995 (27.07.95) (30) Priority Data: 9415167.7 27 July 1994 (27.07.94) GB (71) Applicant (for all designated States except US): CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regents Park, London NW1 4JL (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): SPRINGER, Caroline, Joy [GB/GB]; The Institute of Cancer Research, The Royal Cancer Hospital; 15 Cotswold Road, Belmont, Sutton, Surrey SM2 2NG (GB). MARAIS, Richard [GB/GB]; Chester Beatty Laboratories, The Institute of Cancer Research, 237 Fulham Road, London SW3 6JB (GB). (74) Agents: BRASNETT, Adrian, Hugh et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5LX (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  Published Without international search report and to be republished upon receipt of that report.	
(54) Title: INTRACELLULAR EXPRESSION OF CARBOXYPEPTIDASE G2 IN ENZYME PRODRUG THERAPY			
(57) Abstract  The invention provides a two-component system for gene-directed enzyme prodrug therapy which comprises: (a) a vector which encodes a carboxypeptidase which is expressed within a cell; and (b) a prodrug which can be converted into an active drug by said carboxypeptidase, useful in the treatment of tumours.			

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

INTRACELLULAR EXPRESSION OF CARBOXYPEPTIDASE G2  
IN ENZYME PRODRUG THERAPY

The present invention relates to gene directed enzyme prodrug therapy (GDEPT) and its use in the treatment of disease, including tumours.

A therapeutic approach termed "virus-directed enzyme prodrug therapy" (VDEPT) has been proposed as a method for treating tumour cells in patients using prodrugs. Tumour cells are targeted with a viral vector carrying a gene encoding an enzyme capable of activating a prodrug. The gene may be transcriptionally regulated by tissue specific promoter or enhancer sequences. The viral vector enters tumour cells and expresses the enzyme, in order that a prodrug is converted to an active drug within the tumour cells (Huber et al, Proc. Natl. Acad. Sci. USA (1991) 88, 8039). Alternatively, non-viral methods for the delivery of genes have been used. Such methods include calcium phosphate co-precipitation, microinjection, liposomes, direct DNA uptake, and receptor-mediated DNA transfer. These are reviewed in Morgan & French Anderson, Annu. Rev. Biochem., 1993, 62:191. The term "GDEPT" (gene-directed enzyme prodrug therapy) is used to include both viral and non-viral delivery systems.

A number of different enzymes have been proposed as being suitable for use in GDEPT therapy. In general, the enzymes have been selected to be compatible with prodrugs which are known to be effective against the tumour type being treated. It is also important for the success of GDEPT that the drug is capable of entering the cell.

The enzyme carboxypeptidase G2 (CPG2) has been proposed for use in Antibody-Directed Enzyme Prodrug Therapy (ADEPT) systems, in combination with a number of types of prodrugs, including nitrogen mustard prodrugs. In ADEPT, the prodrug is activated outside the cell or on the cell surface, and thus the prodrug was believed to have no significant ability to enter the cell. Also the prodrug

- 2 -

contains the hydrophilic glutamic acid. It has now surprisingly been found that such prodrugs not only enter the cell but can be activated intracellularly by CPG2. Furthermore, following activation in the cell, the drugs exhibit an effective bystander effect where surrounding cells are also killed by activated drug.

#### Disclosure of the invention.

In order to overcome these problems, the present invention thus provides a two component system for gene-directed enzyme prodrug therapy which comprises: (a) a vector which encodes a carboxypeptidase which is expressed within a cell; and (b) a prodrug which can be converted into an active drug by said carboxypeptidase. The enzyme is preferably a bacterial carboxypeptidase, especially carboxypeptidase CPG2 (SEQ ID No. 1) or *Pseudomonas*  $\gamma$ -glutamylhydrolase EC3.4.22.12 (Levy CC & Goldman P J. Biol. Chem. 242; p2933 (1967)).

The vector may be a RNA or DNA vector. It may be derived from a viral vector, including any suitable vector available in the art for targeting tumour cells.

The invention also provides the system of the invention for use in a method of treatment of a patient, and a method of treating a tumour in a patient in need of treatment which comprises administering to said patient an effective amount of a vector encoding a carboxypeptidase and a prodrug capable of being converted by said enzyme to an active drug.

#### Brief Description of the Drawings

Figure 1. Expression of CPG2\* in NIH3T3 cells (A) Immunoblot analysis. NIH3T3 cells were transfected with MLV $\beta$ plink (lane 1) or with MLVCPG2\* (lane 2). The cells were extracted into buffer A and samples of each were electrophoresed in 10%w/v polyacrylamide gels, transferred to nitrocellulose and probed with a CPG2 specific polyclonal antibody. The position of migration of CPG2\* is shown

- 3 -

and the positions of migration of standard molecular mass markers (in kDa) are shown to the left of the figure. (B) CPG2 enzyme activity assay. The cell extracts from part (A) were analysed by CPG2 enzyme activity assay. Sample numbers correspond to lane numbers from part (A). The enzyme activity is expressed in arbitrary units, relative to buffer controls.

Figure 2. Cytotoxicity assay NIH3T3 cells were transfected with MLV $\beta$ splink (samples 1, 3) or MLVCPG2\* (lanes 2, 4). Forty-two hours after transfection, the cells were treated with CMDA prodrug vehicle as a control (lanes 1, 2) or with CMDA prodrug for 24 hours (lanes 3, 4). After a recovery period, the cells were passaged into fresh tissue culture dishes and after three days, the viability of the cells determined by measuring [methyl-<sup>3</sup>H]thymidine incorporation. The amount of incorporated thymidine is shown in terms of dpm for each sample.

Figure 3A shows the constitutive expression of CPG2\* in NIH3T3 cells.

Figures 3B and 3C show in arbitrary units the enzymatic activity for  $\beta$ -galactosidase (A) and CPG2 (B) expressed in NIH3T3 cell lines #3 (open bar) and #9 (solid bar).

#### Detailed description of the invention.

##### A. Vector systems.

Examples of suitable vector systems include vectors based on the Molony murine leukaemia virus are known (Ram, Z et al, Cancer Research (1993) 53;83-88; Dalton & Treisman, Cell (1992) 68; 597-612). These vectors contain the Murine Leukaemia virus (MLV) enhancer cloned upstream at a  $\beta$ -globin minimal promoter. The  $\beta$ -globin 5' untranslated region up to the initiation ATG is supplied to direct efficient translation of the cloned protein. The initiator ATG straddles an NcoI restriction site and thus can be used to clone a protein coding sequence into the vector. This

- 4 -

vector further contains a polylinker to facilitate cloning, followed by the  $\beta$ -globin 3'-untranslated region and polyadenylation sites. The MLV enhancer is of particular use since it is a strong enhancer and is active in most murine and human cell lines.

Suitable viral vectors further include those which are based upon a retrovirus. Such vectors are widely available in the art. Huber et al (ibid) report the use of amphotropic retroviruses for the transformation of hepatoma, breast, colon or skin cells. Culver et al (Science (1992) 256: 1550-1552) also describe the use of retroviral vectors in GDEPT. Such vectors or vectors derived from such vectors may also be used. Other retroviruses may also be used to make vectors suitable for use in the present invention. Such retroviruses include rous sarcoma virus (RSV). The promoters from such viruses may be used in vectors in a manner analogous to that described above for MLV.

Englehardt et al (Nature Genetics (1993) 4: 27-34) describe the use of adenovirus based vectors in the delivery of the cystic fibrosis transmembrane conductance product (CFTR) into cells, and such adenovirus based vectors may also be used. Vectors utilising adenovirus promoter and other control sequences may be of use in delivering a system according to the invention to cells in the lung, and hence useful in treating lung tumours.

Vectors encoding carboxypeptidase may be made using recombinant DNA techniques known per se in the art. The sequences encoding the enzyme may be constructed by splicing synthetic or recombinant nucleic acid sequences together, or modifying existing sequences by techniques such as site directed mutagenesis. Reference may be made to "Molecular Cloning" by Sambrook et al (1989, Cold Spring Harbor) for discussion of standard recombinant DNA techniques. In general, the vector may be any DNA or RNA vector used in VDEPT or GDEPT therapies.

#### B. Carboxypeptidase.

- 5 -

The carboxypeptidase will usually convert the prodrug into an active drug by removing a protecting group from the prodrug. In most cases, the protecting group will be cleaved as a whole from the prodrug. However, it is also possible for the enzyme to cleave or simply alter part of the protecting group, resulting in a partially cleaved or altered protecting group which is unstable, resulting in spontaneous removal of the remainder of the group.

Carboxypeptidase G2 (CPG2) is disclosed in WO88/07378, and its sequence is also set out below as Seq. ID No. 1. Although the CPG2 with the amino acid sequence of Seq. ID No. 1 is preferred, alterations to the sequence which are amino acid substitutions, deletions or insertions (eg. of about 1, 2, 3, 4, 5, 10 or 20 residues in each case) are also possible. In any event, the alteration will be such that the enzyme retains its ability to convert a prodrug to an active drug at substantially the same rate as the native enzyme. In this context, "substantially the same rate" will desirably be within 1 order of magnitude, and preferably from about 50-fold e.g. about 2-fold less to 2, 5 or 10 fold more.

In addition to specific changes the enzyme may otherwise be altered by truncation, substitution, deletion or insertion as long as the activity of the enzyme is substantially unchanged as defined above. For example, small truncations in the N- and/or C-terminal sequence may occur as a result of the manipulations required to produce a vector in which a nucleic acid sequence encoding the enzyme is linked to a suitable promoter. The activity of the altered enzyme may be measured in model systems such as those described in the examples.

An example of a truncated enzyme is CPG2\* which contains amino acids 23 to 415 of Seq. ID No. 1.

### C. Promoters.

The carboxypeptidase will be expressed in the vector using a promoter capable of being expressed in the cell to which the vector is targeted. The promoter will be operably linked to the sequences

- 6 -

encoding the enzyme and its associated sequences. For example, the c-erbB2 proto-oncogene is expressed in breast tissues at low levels and in a tissue restricted manner. In some tumour states however the expression of this protein is increased, due to enhanced transcriptional activity. Notable examples of this are breast tissue (about 30% of tumours), ovarian (about 20%) and pancreatic tumours (about 50-75%). In such tumours where expression of c-erbB2 is increased due to enhanced transcription or translation, the c-erbB2 promoter may be used to direct expression of proteins in a cell specific manner.

With the GDEPT system of the present invention utilising c-erbB2 promoters to target such tumours, the specificity of GDEPT is increased since transfection of normal cells by a vector with a c-erbB2 promoter will provide only very limited amount of enzyme or none and thus limited activation of prodrug.

The c-erbB-2 promoter has been sequenced to -1500 and may be obtained by reference to Hudson et al, (1990) J. Biol. Chem. 265; 4389-4393. The major start site of transcription has been determined by Ishii et al (1987) Proc. Natl. Acad. Sci. 84; 4374-4378 and Tal et al (1987) Mol. Cell Biol. 7; 2597-2601. This start site is referred to as +1 and this numbering is referred to herein. Translation of c-erbB-2 starts at +178. The promoter has a CAAT box at -75 and a TATA box at -25.

Hollywood and Hurst (1993) EMBO J. 12; 2369-2375 report that in mammary cells, regions of the promoter at -100 and -213 are important for the regulation of transcription. (see also Sarkar et al (1994) J. Biol. Chem. 269; 12285-12289).

In order to achieve expression from a vector utilizing the c-erbB-2 promoter, it is desirable to use the c-erbB-2 promoter region from the CAAT box, and preferably the TATA box, upstream to include sequence elements responsible for specificity of expression in particular tissues, such as those found for mammary cells by Hollywood and Hurst (ibid). The promoter will thus desirably



- 7 -

include at least all the nucleotides upstream (5') of the CAAT box to about the 250th, eg. 300th, 400th, 500th, 600th, 700th, 800th, 900th or further nucleotide 5' to the start of transcription. It is also preferred to include the TATA box. Optionally, the c-erbB-2 sequences downstream of the TATA box to the start of translation at +178 may also be used.

Although the human c-erbB-2 promoter sequence is preferred, modified promoter sequences which are capable of selectively hybridizing to the human sequence may also be used. A promoter sequence capable of selectively hybridizing to the human promoter sequence will be generally at least 70%, preferably at least 80 or 90% and more preferably at least 95% homologous to the promoter region or fragment thereof over a region of at least 20, preferably at least 30, for instance 40, 60 or 100 or more contiguous nucleotides.

In general, those of skill in the art will appreciate that some regions of the promoter such as those at -213 will need to be retained to ensure tumour specificity of expression from the vector whereas other regions of the promoter may be modified or deleted without significant loss of specificity. Thus, modified promoters which are transcriptionally regulated substantially to the same degree as human c-erbB-2 are preferred. The degree of regulation of such candidate promoters can be tested and assessed by those of skill in the art using for example CAT assays in accordance with those described by Hollywood and Hurst (ibid).

"Operably linked" refers to a juxtaposition wherein the promoter and the enzyme-coding sequence are in a relationship permitting the coding sequence to be expressed under the control of the promoter. Thus, there may be elements such as 5' non-coding sequence between the promoter and coding sequence which is not native to either the promoter nor the coding sequence. Such sequences can be included in the vector if they do not impair the correct control of the coding sequence by the promoter.

- 8 -

Other suitable promoters include viral promoters such as mammalian retrovirus or DNA virus promoters. Suitable promoters include those used in vectors described above, e.g. MLV, CMV, RSV and adenovirus promoters. Preferred adenovirus promoters are the adenovirus early gene promoters. Strong mammalian promoters may also be suitable. An example of such a promoter is the EF-1 $\alpha$  promoter which may be obtained by reference to Mizushima and Nagata ((1990), Nucl. Acids Res. 18; 5322). Variants of such promoters retaining substantially similar transcriptional activities may also be used.

The invention provides a viral vector comprising a c-erbB-2 promoter operably linked to a gene encoding a carboxypeptidase enzyme, the enzyme being capable of converting a prodrug into an active drug. The invention also provides a kit which comprises a vector as defined above together with a prodrug which is capable of being converted to an active drug by the enzyme encoded by the vector. In another aspect, the invention provides a vector as defined above or a kit as defined above for use in a method of treatment of the human or animal body, and in particular a method of treatment of tumours in which the c-erbB-2 proto-oncogene is overexpressed. In a further aspect, the invention provides a method of treatment of tumours, and in particular a method of treatment of tumours in which the c-erbB-2 proto-oncogene is overexpressed, which method comprises administering to an individual with a tumour (i) an effective amount of a vector as defined above, and (ii) an effective amount of a prodrug capable of being converted to an active drug by the enzyme encoded by the vector.

#### D. Prodrugs.

The prodrug for use in the system will be selected to be compatible with the enzyme, ie. such that the enzyme will be capable of converting the prodrug into an active drug. Desirably, the toxicity of the prodrug to the patient being treated will be at least one order of magnitude less toxic to the patient than the active drug. Preferably, the active drug will be several, eg 2,

- 9 -

3, 4 or more orders of magnitude more toxic. Suitable prodrugs include nitrogen mustard prodrugs and other compounds such as those described in WO88/07378, WO89/10140, WO90/02729, WO91/03460, EP-A-540 263, WO94/02450, WO95/02420 or WO95/03830 which are incorporated herein by reference.

E(1) - Nitrogen mustard prodrugs.

Nitrogen mustard prodrugs include compounds of the formula:




where Ar represents an optionally substituted ring aromatic ring system, R-NH is the residue of an  $\alpha$ -amino acid R-NH<sub>2</sub> or oligopeptide R-NH<sub>2</sub> and contains at least one carboxylic acid group, and M represents a nitrogen mustard group.

The residue of the amino acid R-NH is preferably the residue of glutamic acid. It is disclosed in WO88/07378 that the enzyme carboxypeptidase G2 is capable of removing the glutamic acid moiety from compounds of the type shown above, and the removal of the glutamic acid moiety results in the production of an active nitrogen mustard drug.

- 10 -

Thus nitrogen mustard prodrugs of use in the invention include the prodrugs of generic formula I of WO94/02450 and salts thereof, and in particular those of formula (I):



wherein  $R^1$  and  $R^2$  each independently represent chlorine, bromine, iodine,  $\text{OSO}_2\text{Me}$ ,  $\text{OSO}_2\text{phenyl}$  (wherein phenyl is optionally substituted with 1,2,3,4 or 5 substituents independently selected from  $\text{C}_{1-4}$  alkyl, halogen,  $-\text{CN}$  or  $\text{NO}_2$ );

$R^{1a}$  and  $R^{2a}$  each independently represents hydrogen,  $\text{C}_{1-4}$  alkyl or  $\text{C}_{1-4}$  haloalkyl;

$R^3$  and  $R^4$  each independently represents hydrogen,  $\text{C}_{1-4}$  alkyl or  $\text{C}_{1-4}$  haloalkyl;

$n$  is an integer from 0 to 4;

each  $R^5$  independently represents hydrogen,  $\text{C}_{1-4}$  alkyl optionally containing one double bond or one triple bond,  $\text{C}_{1-4}$  alkoxy, halogen, cyano,  $-\text{NH}_2$ ,  $-\text{CONR}^7\text{R}^8$  (wherein  $R^7$  and  $R^8$  are independently hydrogen,  $\text{C}_{1-6}$  alkyl or  $\text{C}_{3-6}$  cycloalkyl) or two adjacent  $R^5$  groups together represent

a)  $\text{C}_4$  alkylene optionally having one double bond;

b)  $\text{C}_3$  alkylene; or

c)  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ ,  $-\text{CH}-\text{CH}-\text{CH}_2-$  or  $-\text{CH}_2-\text{CH}=\text{CH}-$  each optionally substituted with 1, 2, 3 or 4 substituents said substituents each independently selected from the group consisting of  $\text{C}_{1-4}$  alkyl,  $\text{C}_{1-4}$  alkoxy, halogen, cyano and nitro;

$X$  is a group  $-\text{C}(\text{O})-$ ,  $-\text{O}-\text{C}(\text{O})-$ ,  $-\text{NH}-\text{C}(\text{O})-$  or  $-\text{CH}_2-\text{C}(\text{O})-$ ; and

$Z$  is a group  $-\text{CH}_2-\text{T}-\text{C}(\text{O})-\text{OR}^6$  where  $T$  is  $\text{CH}_2$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-(\text{SO})-$  or  $-(\text{SO}_2)-$ , and  $R^6$  is hydrogen,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{3-6}$  cycloalkyl

- 11 -

amino, mono- di- $C_{1-6}$  alkylamino or mono or di $C_{3-6}$  cycloalkyl amino, provided that when  $R^6$  is hydrogen T is  $-CH_2-$ ; and physiologically acceptable derivatives, including salts, of the compounds of formula (I).

Halogen includes fluorine, chlorine, bromine and iodine. Preferred values for the groups  $R^{1a}$  and  $R^{2a}$  are methyl and hydrogen, especially hydrogen. Preferred values for the groups  $R^3$  and  $R^4$  are hydrogen, methyl and trifluoromethyl, especially hydrogen. Preferred values for the groups  $R^1$  and  $R^2$  are I, Br, Cl,  $OSO_2Me$  and  $OSO_2phenyl$  wherein phenyl is substituted with one or two substituents in the 2 and/or 4 positions. I, Cl and  $OSO_2Me$  are especially preferred.

Preferred values for  $R^5$  when n is an integer from 1 to 4 are fluorine, chlorine, methyl- $CONH_2$  and cyano. Preferably, n is 0, 1 or 2. When n is 1 or 2 it is preferred that  $R^5$  is fluorine at the 3 and/or 5 positions of the ring. The group X is preferably  $-C(O)-$ ,  $-O-C(O)-$  or  $-NH-C(O)-$ . Z is preferably a group  $-CH_2CH_2-COOH$ .

Preferred specific compounds include:

N-4-[(2-chloroethyl)(2-mesyloxyethyl)amino]benzoyl-L-glutamic acid (referred to below as "CMDA") and salts thereof;

N-(4-[bis(2-chloroethyl)amino]-3-fluorophenylcarbamoyl)-L-glutamic acid and salts thereof;

N-(4-[bis(2-chloroethyl)amino]phenylcarbamoyl)-L-glutamic acid and salts thereof;


N-(4-[bis(2-chloroethyl)amino]phenoxy-carbonyl)-L-glutamic acid and salts thereof; and

N-(4-[bis(2-iodoethyl)amino]phenoxy-carbonyl)-L-glutamic acid and salts thereof.

Particular sub-groups of the compounds of the present invention of interest may be obtained by taking any one of the above mentioned particular or generic definitions for  $R^1-R^4$ ,  $R^5$ , X or W either singly or in combination with any other particular or generic definition for  $R^1-R^4$ ,  $R^5$ , X or W.

- 12 -

Prodrugs which are more lipophilic than compounds of the prior art such as those disclosed in WO88/07378 are of use in the system of the invention, including compounds of the formula (II):



wherein  $R^1$ ,  $R^2$ ,  $R^5$ ,  $n$  and  $Z$  are as defined for compounds of the formula (I) above;

$m$  is an integer from 0 to 4,

$Z^1$  and  $Z^2$  are each independently -O- or -NH-; and

$R^9$  is hydrogen, *t*-butyl or allyl;

and physiologically acceptable derivatives of the compound of formula (I). Preferred values of  $R^1$ ,  $R^2$ ,  $R^5$ ,  $n$  and  $Z$  are as defined above for compounds of the formula (I). Preferred values of  $m$  are 0, 1 or 2 as defined for  $n$  above.  $R^9$  is preferably hydrogen, but can be protected especially during synthesis by groups such as allyl or *t*-butyl.


These prodrugs can be activated at the site of a tumour by a carboxypeptidase enzyme, for example, CPG2 as disclosed in WO88/07378 or WO94/02450.

Compounds of the formula (II) may be made using reactions and methods known per se in the art of chemistry. The following methods are of particular use:


A: Compounds of formula (II) where  $Z^1$  is -O-:

- 13 -

Compounds of the formula (I) in which  $Z^1$  is -O- may be made by reacting a nitrogen mustard of formula (III)



where  $R^1$ ,  $R^2$ , and  $R^5$  and  $n$  are as defined above and  $Z^4$  is -O- with a linker of formula (IV)




where  $R^5$ ,  $m$ ,  $Z^2$ ,  $R^9$  and  $Z^3$  are as defined above, and  $Q$  is hydrogen or a leaving group. This reaction may be done in aprotic solvents in the presence of a base, for example DMF and triethylamine.

Preferred leaving groups  $Q$  include a succinimidyl group, a 4-nitrophenyl carbonate group, a pentafluorophenyl carbonate and a tetrachloroethyl group  $CH(Cl)CCl_3$ .

- 14 -

(ii) Compounds of the formula (III) may be made starting from 4-nitrophenol optionally substituted with the group(s)  $R^5_{(n)}$  (as defined above). The phenolic group is protected as an adamantanyloxycarbonyl-derivative (by reacting the starting materials with adamantanyl-fluoroformate and triethylamine in THF at room temperature). The protected 4-nitrophenyl carbonate is reduced to the corresponding amine by hydrogen transfer in ethanol using ammonium formate and Pd/C 10% as catalyst at rt. The amine is then hydroxyethylated with ethylene oxide in AcOH at 20°C and then reacted to the desired nitrogen mustard. Reference may be made to EP-A-433 360 or EP-A-490970 for suitable conditions. The compounds may be purified by column chromatography. Deprotection to remove the adamantyl group may be carried out in trifluoroacetic acid.

(iii) Alternatively, the nitrogen mustard of formula (III) may be activated as a chloroformate by treatment with phosgene or triphosgene in an aprotic solvent and triethylamine followed by coupling with a compound of formula (V):



where  $R^5$ ,  $m$ ,  $Z^2$ ,  $R^9$  and  $Z^1$  are as defined above. This may be carried out in THF or other aprotic solvents in the presence of a base (for example triethylamine or pyridine).



- 15 -

(iv) A further alternative route of synthesis of compounds of the formula (II) in which  $Z^1$  is -O- involves direct coupling of 4-nitrophenol optionally substituted with the group(s)  $R^5_{(n)}$  (as defined above) with the compound of the formula (IV) or by reaction of the said optionally substituted 4-nitrophenol compound chloroformate with the compound of formula (IV), followed in each case by the reaction described above to convert the nitro group, via an amine, to a mustard group.

B: Compounds of formula (II) where  $Z^1$  is -NH-:


(i) Compounds of the formula (II) in which  $Z^1$  is -NH- may be made by reaction of a compound of formula (III) in which  $Z^4$  is -NH- with a linker of the formula (IV) in aprotic solvents and in the presence of a base. Compounds of the formula (III) in which  $Z^4$  is -NH- may be made from a 1-halo-4-nitrobenzy compound, optionally substituted with the group(s)  $R^5_{(n)}$  (as defined above). This is converted to the corresponding 1-bis-hydroxyethylamino-4-nitrobenzyl compound by reaction with diethanolamine with heat and the resulting product purified by column chromatography. The corresponding 4-nitro nitrogen mustard may be made by for example mesylation using mesyl chloride in pyridine and subsequent reaction to other halo mustards, e.g. bromo or iodo mustards if required. The 4-nitro group may be reduced by hydrogen transfer in ethanol using ammonium formate and a Pd/C 10% catalyst at 20°C.

(ii) Alternatively the 1-bis-hydroxyethylamino-4-nitrobenzyl compound mentioned above can be reduced using ammonium formate and Pd/C 10% as catalyst in ethanol at 20°C to provide the corresponding phenylene-diamino derivative. This derivative can be converted into the corresponding 4-amino nitrogen mustard as described in the above paragraph, e.g. initially by reaction with mesyl chloride.

C: Compounds of the formula (IV) in which  $Z^2$  is -NH-:

- 16 -

(i) Compounds of the formula (IV) in which  $Z^2$  is  $-NH-$  may be made from a 4-nitro benzylic alcohol optionally substituted with the group(s)  $R^5(m)$  (as defined above). The hydroxyl function is protected as a pyranyl- or *t*-butyl-dimethylsilyl (TBDMSi)-ether by treatment at 20°C with 3,4-2H-dihydropyran and pyridinium-*p*-toluensulfonate (PPTS) in an aprotic solvent or with TBDMSi chloride and imidazole in dimethylformamide (DMAC), respectively. The intermediate thus obtained is reduced to the corresponding amine by hydrogen transfer in ethanol using ammonium formate and Pd/C 10% as catalyst at 20°C. This amine is converted to a glutamyl ester intermediate of formula (VI):



(VI)

where  $R^5$ ,  $m$ ,  $R^9$  and  $Z^1$  are as defined above,  $Z^2$  is  $-NH-$  and Pr is the pyranyl- or *t*-butyl-dimethylsilyl (TBDMSi)-ether protecting group. This may be done by treating the amine with triphosgene and triethylamine in toluene at 60°C to provide the corresponding isocyanate, which is treated with a glutamate derivative of formula  $R^9-C(O)-CH(NH_2)-Z^1$  where  $R^9$  and  $Z$  are as defined above. Alternatively the corresponding glutamyl-isocyanate obtained from the corresponding glutamate by treatment with triphosgene and triethylamine in toluene at -78°C may be reacted with the amine in a one pot procedure.

(ii) The compound of formula (VI) is deprotected to remove the TBDMSi or pyranyl groups by treatment with mild acidic media (AcOH,

- 17 -

THF and H<sub>2</sub>O or PPTS, EtOH, 55°C). This yields a compound of formula (VI) in which Pr is hydrogen. Compounds of the formula (IV) in which Q is a leaving group may be prepared using standard reactions known in the art.

(iii) For example where Q is a succinimidyl group the compound of formula (VI) where Pr is hydrogen may be treated with disuccinimidyl-carbonate and triethylamine in acetonitrile. Where a 4-nitrophenyl carbonate group is desired treatment with 4-nitrophenyl chlorformate and triethylamine in THF may be used. A pentafluorophenyl carbonate may be added by in situ phosgenation of pentafluorophenol followed by coupling to the linker of formula (VI) in which Pr is hydrogen.

D: Compounds of the formula (IV) in which Z<sup>2</sup> is -O-:

(i) The starting materials for the linkers possessing a carbamic bond are unsubstituted or substituted (with the group(s) R<sup>5</sup><sub>(n)</sub> (as defined above)) 4-hydroxy-benzylic alcohols. These type of linkers may require an extra electron withdrawing group on the aromatic nucleus in order to undergo 1,4-elimination. The 4-hydroxy group is specifically protected as an acetate by treating the starting material with acetyl-v-triazolo-[4,5-b]pyridine, 1N NaOH in THF at 20°C. The alcohol function of the acetate is further protected as pyranyl- or TBDMSi-ether by the procedures described in section C above. The acetate function is then deprotected to restore the 4-hydroxy group in NaHCO<sub>3</sub> aq. MeOH at 20°C. The resulting phenol compounds are reacted in a one pot procedure with a protected glutamyl-isocyanate as described in section C(i) above. This yields a compound of the formula of (VII) as shown above in which Z<sup>2</sup> is -O- and Pr is the pyranyl- or t-butyl-dimethylsilyl (TBDMSi)-ether protecting group.

(ii) Deprotection of this compound yields a compound of the formula (VI) in which Pr is hydrogen. This may be converted to compounds of the formula (IV) by methods analogous to those described in sections C(ii) and (iii) above.

- 18 -



E: Alternative synthesis of compounds of formula (III):

Compounds of the formula (III) in which Q is hydrogen, fluoro, chloro, bromo or -O-(N-succinimide) may also be obtained by reference to WO95/02420 or WO95/03830.

E(ii). Other prodrugs.

Further prodrugs suitable for use in the invention include those of the general formula: FTLi-(PRT)<sub>m</sub>, or salts thereof where FTLi is a ras inhibitor such as a farnesyltransferase inhibitor compound and PRT represents m' protecting groups capable of being cleaved from the ras inhibitor by the action of an enzyme, where m' is an integer from 1 to 5. Such compounds are disclosed in WO95/03830.

Suitable FTLi's include those of the formula



where R<sub>3a</sub> and R<sub>4a</sub> are the side chains of naturally occurring amino acids (for example -CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>, -CH(OH)CH<sub>3</sub>), including their oxidized forms (for example, methionine sulfoxide or methionine sulfone), or are substituted or unsubstituted aliphatic, aromatic or heteroaromatic groups, preferably cyclohexyl, phenyl, pyridyl, imidazolyl or saturated or unsaturated branched or straight chains of 2 to 8

- 19 -

carbon atoms optionally substituted with an aromatic or heteroaromatic ring;


$X_{1a}$  is  $CH_2CH_2$ , trans  $CH=CH$  or  $CH_2NH$ ; and

$X_{2a}$  is  $(CH_2)_n$  wherein  $n$  is 0, 1 or 2. Preferably  $R_{3a}$  and  $R_{4a}$  both represent  $CH(CH_3)CH_2CH_3$ .

Suitable FLT<sub>i</sub> compounds including those of the above formulae may be obtained by reference to WO95/03830.

A further group of prodrugs are tyrphostin compounds of the general formula: PTKi-PRT<sub>m</sub>, where PTKi is a compound with PTK (protein tyrosine kinase) inhibitory activity, PRT is a protecting group capable of being cleaved from the PTK inhibitor by the action of an enzyme and  $m'$  is an integer from 1 to 5.

Suitable PTKs include tyrphostins. Tyrphostins are low molecular weight (e.g. less than 2,000) styryl containing inhibitors of protein tyrosine kinase which are capable of binding to the subsite of protein tyrosine kinase domains. Suitable tyrphostins include those described by Gazit et al (Gazit et al, J. Med. Chem. (1989) 32, 2344) and Gazit et al (J. Med. Chem. (1991) 43; 1896-1907) and especially tyrphostins of the general formula:



where  $X^1$  represents carbon, a nitrogen or a group  $N=O$ ;  $n$  is an integer from 1 to 3; each group  $R^{10}$ , which may be the same or different is hydrogen, hydroxy, mercapto, carboxy, formyl,  $C_{1-4}$ alkyl,  $C_{2-4}$  alkenyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkylthio, carboxy $C_{1-4}$ alkyl, carboxy $C_{2-4}$  alkenyl,  $C_{1-4}$ alkylsulphoxy, halo (ie. fluoro, chloro, bromo or iodo), nitro, amino,  $C_{1-4}$ alkylamino, or  $C_{1-4}$ dialkylamino, or when  $n$  is 2 or 3 two  $R^{10}$  groups may together form a methylenedioxy or ethylenedioxy group;  $R^{11}$  is hydrogen, hydroxy,  $C_{1-4}$ alkyl or

- 20 -

together with position 2 of the ring to which the group(s)  $R^{10}$  is(are) attached forms a 5 or 6 membered aliphatic or heterocyclic ring, said 5 or 6 membered ring optionally containing a ketone group; and  $R^{12}$  is cyano, carboxy, carbamoyl, thiocarbamoyl, a group  $C(O)HNCH_2CN$ , a group  $C(NH_2)=C(CN)_2$ , an alpha keto group  $C(O)R^{13}$  where  $R^{13}$  is 3,4-dihydroxyphenyl or 2-thiophenyl or an alpha amido group  $C(O)NHR^{14}$  where  $R^{14}$  is benzyl, phenyl or 2,4-dimethoxyphenyl; provided that at least one of the groups  $R^{10}$  and  $R^{11}$  is mercapto, hydroxy or amino.


In a preferred embodiment,  $x^1$  is C; n is an integer from 1 to 3; each group  $R^{10}$ , which may be the same or different is hydrogen, hydroxy, carboxy, formyl,  $C_{1-4}$ alkyl,  $C_{2-4}$  alkenyl,  $C_{1-4}$ alkoxy, carboxy $C_{1-4}$ alkyl, carboxy $C_{2-4}$  alkenyl, halo (ie. fluoro, chloro, bromo or iodo), nitro, amino,  $C_{1-4}$ alkylamino, or  $C_{1-4}$ dialkylamino, or when n is 2 or 3 two  $R^{10}$  groups may together form a methylenedioxy or ethylenedioxy group;  $R^{11}$  is hydrogen, hydroxy or  $C_{1-4}$ alkyl; and  $R^{12}$  is cyano, carboxy, carbamoyl, thiocarbamoyl, a group  $C(O)HNCH_2CN$  or a group  $C(NH_2)=C(CN)_2$ . Most preferably,  $x^{10}$  represents carbon, n is an integer from 1 to 3; each group  $R^{10}$ , which may be the same or different is hydrogen, hydroxy or amino;  $R^{11}$  is hydrogen or hydroxy; and  $R^{12}$  is cyano, a group  $C(O)HNCH_2CN$ , a group  $C(NH_2)=C(CN)_2$ , an alpha keto group  $C(O)R^{13}$  where  $R^{13}$  is 3,4-dihydroxyphenyl, or an alpha amido group  $C(O)NHR^{14}$  where  $R^{14}$  is benzyl; provided that at least one of the groups  $R^{10}$ ,  $R^{11}$ , and  $R^{12}$  are hydroxy or amino. Preferably,  $R^{10}$  is hydroxy or amino. When  $R^{11}$  forms a 5 or 6 membered ring with  $R^{10}$  preferred rings include heterocyclic rings wherein the ring contain one nitrogen atom and 4 or 5 carbon atoms. The total number of atoms includes the 2 carbon atoms of the ring to which the group(s)  $R^{10}$  is(are) attached.

Suitable tyrphostins such as the above may be obtained by the methods disclosed in, or analogous to those of, WO95/02420, Gazit et al 1989 and 1991, *ibid*, which are incorporated herein by reference.

Tyrphostins may be linked to any suitable protecting group which is removable by an enzyme. Suitable protecting groups PRT may be

- 21 -

found by reference to WO95/03830 or WO95/02420 and may be of the structure:



where  $Z^1$  is as defined above or S,  $m'$  is an integer from 1 to 5, Ph is a phenylene ring optionally substituted by from 1 to 4 groups  $R^5$  (which may be the same or different) as defined above and glu is a group  $-CHZ-C(O)-OR^9$  where Z and  $R^9$  are as defined above.

#### E(iii). Derivatives.

Physiologically acceptable derivatives of said prodrug include salts, amides, esters and salts of esters. Esters include carboxylic acid esters in which the non-carbonyl moiety of the ester grouping is selected from straight or branched chain  $C_{1-6}$  alkyl, (methyl, n-propyl, n-butyl or t-butyl); or  $C_{3-6}$  cyclic alkyl (e.g. cyclohexyl). Salts include physiologically acceptable base salts, eg derived from an appropriate base, such as alkali metal (e.g. sodium), alkaline earth metal (e.g. magnesium) salts, ammonium and  $NR_4^+$  (wherein  $R^n$  is  $C_{1-4}$  alkyl) salts. Other salts include acid addition salts, including the hydrochloride and acetate salts. Amides include non-substituted and mono- and di-substituted derivatives.

#### F. Applications of the invention

The system of the invention can be used in a method of treatment of the human or animal body. Such treatment includes a method of treating the growth of neoplastic cells which comprises administering to a patient in need of treatment the system of the invention. It is also possible that the invention may be used to treat cells which are diseased through infection of the human or animal body by bacteria, viruses or parasites. Viral late promoters often rely on viral proteins that are made early in the infection. The viral coat proteins which are expressed on the surface of an infected cell may be used as a target for getting the gene into the cell. If a viral late promoter is then used to direct expression of the GDEPT enzyme, any infected cells will express the protein, and specifically, cells which have been infected, for some time. This may be sufficient to kill the infected cells. For parasites, a parasitic promoter and parasite

- 22 -

surface proteins, may be used to direct expression and infect the parasites respectively.

For a bacteria, all the delivery systems will probably need to be changed to use bacterial viruses, although a specific promoter should be easier to define.

For use of the vectors in therapy, the vectors will usually be packaged into viral particles and the particles delivered to the site of the tumour, as described in for example Ram et al (ibid). The viral particles may be modified to include an antibody, fragment thereof (including a single chain) or tumour-directed ligand to enhance targetry of the tumour. Alternatively the vectors may be packaged into liposomes. The liposomes may be targeted to a particular tumour. This can be achieved by attaching a tumour-directed antibody to the liposome. Viral particles may also be incorporated into liposomes. The particles may be delivered to the tumour by any suitable means at the disposal of the physician. Preferably, the viral particles will be capable of selectively infecting the tumour cells. By "selectively infecting" it is meant that the viral particles will primarily infect tumour cells and that the proportion of non-tumour cells infected is such that the damage to non-tumour cells by administration of a prodrug will be acceptably low, given the nature of the disease being treated. Ultimately, this will be determined by the physician.

Suitable liposomes include, for example, those comprising the positively charged lipid (N[1-(2,3-dioleyloxy)propyl]-N,N,N-triethylammonium (DOTMA), those comprising dioleoylphosphatidylethanolamine (DOPE), and those comprising 3 $\beta$ [N-(n',N'-dimethylaminoethane)-carbamoyl]cholesterol (DC-Chol).

One suitable route of administration is by injection of the particles in a sterile solution. While it is possible for the prodrugs to be administered alone it is preferable to present them as pharmaceutical formulations. The formulations comprise a prodrug, together with one or more acceptable carriers thereof and optionally other therapeutic ingredients. The carrier or carriers must be "acceptable" in the sense of being compatible with the



- 23 -

other ingredients of the formulation and not deleterious to the recipients thereof, for example, liposomes. Suitable liposomes include, for example, those comprising the positively charged lipid (N[1-(2,3-dioleyloxy)propyl]-N,N,N-triethylammonium (DOTMA), those comprising dioleoylphosphatidylethanolamine (DOPE), and those comprising 3 $\beta$ [N-(n',N'-dimethylaminoethane)-carbamoyl]cholesterol (DC-Chol).

Viruses, for example isolated from packaging cell lines may also be administered by regional perfusion or direct intratumoral direction, or direct injection into a body cavity (intracavitary administration), for example by intra-peritoneum injection.

It is also known that muscle cells can take up naked DNA and thus sarcomas may be treated using a vector system of the invention in which naked DNA is directly injected into the sarcoma.

Formulations suitable for parenteral or intramuscular administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats, bactericidal antibiotics and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents, and liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water, for injections, immediately prior to use. Injection solutions and suspensions may be prepared extemporaneously from sterile powders, granules and tablets of the kind previously described.

It should be understood that in addition to the ingredients particularly mentioned above the formulations may include other agents conventional in the art having regard to the type of formulation in question. Of the possible formulations, sterile pyrogen-free aqueous and non-aqueous solutions are preferred.

- 24 -

The doses may be administered sequentially, eg. at daily, weekly or monthly intervals, or in response to a specific need of the patient.

Preferred routes of administration are oral delivery and injection, typically parenteral or intramuscular injection or intratumoural injection.

In using the system of the present invention the prodrug will usually be administered following administration of the vector encoding an enzyme. Typically, the vector will be administered to the patient and then the uptake of the vector by transfected or infected (in the case of viral vectors) cells monitored, for example by recovery and analysis of a biopsy sample of targeted tissue.

The exact dosage regime will, of course, need to be determined by individual clinicians for individual patients and this, in turn, will be controlled by the exact nature of the prodrug and the cytotoxic agent to be released from the prodrug but some general guidance can be given. Chemotherapy of this type will normally involve parenteral administration of both the prodrug and modified virus and administration by the intravenous route is frequently found to be the most practical. For glioblastoma the route is often intratumoural. A typical dosage range of prodrug generally will be in the range of from about 1 to 150 mg per kg per patient per day, which may be administered in single or multiple doses. Preferably the dose range will be in the range from about 10 to 75, e.g. from about 10 to 40, mg per kg per patient per day. Other doses may be used according to the condition of the patient and other factors at the discretion of the physician.

Tumours which may be treated using the system of the present invention include any tumours capable of being treated by a GDEPT or VDEPT system and thus are not limited to any one particular class of tumours. Particularly suitable tumour types include breast, colorectal and ovarian tumours, as well as pancreatic, melanoma, glioblastoma, hepatoma, small cell lung, non-small cell lung, muscle and prostate tumours.

- 25 -

The system of the invention may also be used to treat infectious diseases, for example, and any other condition which requires eradication of a population of cells.

It will be understood that where treatment of tumours is concerned, treatment includes any measure taken by the physician to alleviate the effect of the tumour on a patient. Thus, although complete remission of the tumour is a desirable goal, effective treatment will also include any measures capable of achieving partial remission of the tumour as well as a slowing down in the rate of growth of a tumour including metastases. Such measures can be effective in prolonging and/or enhancing the quality of life and relieving the symptoms of the disease.

The following Examples illustrate the invention.

#### PREPARATIVE EXAMPLE 1

The native CPG2 protein contains a signal peptide which serves to export it out of bacterial cells into the extracellular environment. In order to ensure that the signal peptide did not direct the export of CPG2 from mammalian cells, PCR directed site specific mutagenesis was used to delete the signal peptide from the CPG2 gene. This was achieved using the PCR primer #4476:

5'>CGC GGA TCC GCC CTG GCC CAG AAG CGC<3'

in conjunction with primer #4477:

5'>CGC GAA TTC CTT GCC GGC GCC CAG ATC<3'

Primer #4476 was used to alter codons 21 and 22 of the CPG2 gene into a BamH1 restriction endonuclease site. Primer #4477 was used to change codon 416 (the natural stop codon of CPG2) into an EcoR1 restriction endonuclease site. A PCR generated fragment, containing the entire CPG2 open reading frame was generated and digested with BamH1 and EcoR1 and cloned into those sites of the plasmid MLV $\beta$ plink (Dalton and Treisman (1992), Cell 68, 597-612). This results in a CPG2 protein which has the structure: Met-Ala-Gly-Ser-(CPG2 codons 23 to 415)-Glu-Phe-Leu-Glu-Ile-Asp, and therefore lacks the signal peptide. The plasmid is referred to as "MLVCPG2\*" and the protein produced from it as CPG2\*.

- 26 -

NIH3T3 cells were transfected with MLVCPG2\*, or with MLV $\beta$ spink as a control in order to determine whether CPG2 could be produced in mammalian cells. Transfection was performed using the lipid reagent lipofectAMINE (GibcoBRL). For transfection,  $1 \times 10^5$  cells were plated into 35mm tissue culture dishes and incubated at 37°C overnight. The following day, lipofectAMINE:DNA complexes were prepared by combining 0.4 $\mu$ g of plasmid with 6 $\mu$ l of lipofectAMINE reagent (Gibco/BRL) in a total of 32 $\mu$ l of PBSA; the complexes were incubated at room temperature for 15 minutes. The cells were washed twice with 2 ml of serum free DMEM medium and placed in 0.8ml of serum free DMEM. The lipofectAMINE:DNA complexes were diluted into 0.2ml of serum free medium and added to the cells. The cells were incubated with the lipofectAMINE:DNA complexes for 6 hours at 37°C, washed twice with 2ml of DMEM medium containing 10% foetal calf serum (FCS) and placed into 2.5ml 10%FCS/DMEM. After a further incubation period of 42 hours, the cells were washed twice with 5ml PBSA and extracted on the tissue culture dishes into 50 $\mu$ l of buffer A (250mM Tris.HCl, 10%v/v glycerol, 1%v/v Triton X-100, pH7.5). The extract was transferred to 1.5ml tubes, centrifuged at 13,000 rpm in a micro-fuge and the supernatants collected; the protein concentration was determined with the Bio-Rad protein assay kit, with BSA as a standard.

For protein-immunoblot analysis, 10 $\mu$ l of each extract was combined with 10 $\mu$ l of SDS sample buffer and the proteins were separated on a 10 % SDS-polyacrylamide gel (Laemmli U.K. (1970) Nature 227: 680-685). The proteins in the gel were transferred to nitrocellulose by electroelution and the cell extracts analysed by protein immuno-blotting (Gershoni J.M. and Palade G.E. (1983) Anal. Biochemistry 131:1-15), using a CPG2 specific rabbit polyclonal antibody which was raised to CPG2 expressed in Sf9 insect cells. The antibody was used at a dilution of 1:1000 and the immunocomplexes detected using the ECL immunoblotting kit, according to the manufacturers instructions (Amersham).

The extracts were also tested for CPG2 enzyme activity by measuring their ability to cleave the CMDA prodrug into active drug, which can be detected by measuring the change in optical density absorbance at 305nm. For the assay, 5 $\mu$ g of NIH3T3 protein

- 27 -

extracted in buffer A was incubated in 1ml assay buffer (20mM Tris.HCl, 1mM MgCl<sub>2</sub>, 60μM ZnCl<sub>2</sub>, pH 7.4) containing 50μM CMDA prodrug, at 37°C for 15 min. After incubation, EDTA was added to a final concentration of 10mM and the change in absorbance at 305nm was monitored and compared to that of the starting solution. A reduction in absorbance indicates conversion of the prodrug into drug and therefore by inference demonstrates the presence of CPG2; the enzyme activity is expressed in arbitrary units, compared to extraction buffer controls.

From the immunoblot analysis, it could be seen that extracts from cells transfected with MLVCPG2\* contained a protein with an apparent Mr of ~40,000 which was recognised by the CPG2 antibody and which was not present in cells transfected with MLVβplink (figure 1A). When the enzymatic activity of these extracts was examined, CPG2 activity was detected in cell extracts from cells transfected with MLVCPG2\* but not in extracts from cells transfected with MLVβplink (Figure 1B). These data show that it is possible to express CPG2\* inside mammalian cells in a form that is enzymatically active.

#### EXAMPLE 1

The internally expressed CPG2\* was tested for its ability to direct cell death in NIH3T3 cells treated with prodrugs. Two sets of cells were prepared, each of which contained one dish of cells transfected with MLVβplink and one dish of cells transfected with MLVCPG2\*. The cells were incubated for 42 hours following the transfection and then set 1 was incubated in the presence of prodrug vehicle (5mM Hepes, 0.5%v/v DMSO, pH7.0; final concentrations) for 24 hours and set 2 were incubated with the CMDA prodrug (0.5mM final concentration) for 24 hours. Following prodrug treatment, the cells were washed twice with 5ml fresh medium and incubated for a further 24 hours. They were then passaged into fresh culture dishes, seeded at 1x10<sup>5</sup> cells/ 35mm dish in 2ml 10%FCS/ DMEM. After a further 72 hour incubation cell growth was assessed by thymidine incorporation. For thymidine incorporation, 1μCi of [methyl-<sup>3</sup>H]thymidine was added to each well (2ml medium) and incubated for 5 hours at 37°C. The cells were washed twice with PBSA (4°C) and fixed with 5%w/v trichloro acetic

- 28 -

acid (TCA) in water at 4°C for 20 minutes. The TCA was removed and the cells washed twice with 2ml methanol and air dried. The DNA was extracted with 1ml 0.1M NaOH, 1% SDS at room temperature for 5 minutes, added to 4 ml scintillation fluid and the incorporated thymidine was determined by scintillation counting.

Cells transfected with MLVCPG2\* grew at the same rate as cells transfected with MLV $\beta$ plink, indicating that expression of CPG2\* in cells did not affect the growth of these cells (Figure 2, compare samples 1,2). Treatment of the control cells with CMDA had only a minimal effect on their growth rate (Figure 2, compare samples 1,3). By contrast, treatment of the cells expressing CPG2\* with CMDA resulted in completed inhibition of thymidine incorporation and cell death (Figure 2, compare samples 2,4). Thus, these data show that the enzyme CPG2 can be expressed in NIH3T3 cells in an active form which does not appear to affect the rate of growth of these cells. If cells expressing CPG2\* are treated with 0.5mM CMDA for 24 hours, 100% of the cells are killed.

## PREPARATIVE EXAMPLE 2

### 1. Construction of the vector

An expression cassette was constructed by fusing the 1.2kb HindIII/EcoRI fragment from the plasmid pEF-Bos (Mizushima and Nasgata (1990) NAR 18, 5322), which contains the EF1 $\alpha$  promoter, first exon, first intron and a portion of the second exon to the  $\beta$ -globin gene at position -4 relative to the start of transcription of the  $\beta$ -globin gene. The Nco-I/HindIII fragment from the plasmid MLV $\beta$ -Plink, containing the polylinker and the  $\beta$ -globin 3' untranslated region was fused to the Nco-I site of the  $\beta$ -globin gene, which is located across the  $\beta$ -globin start of translation (position +50, relative to the start of transcription), to provide a polylinker and 3' untranslated region for proper mRNA processing.

This entire cassette was to be cloned into the vector pMC1Neo PolyA (Stratagene). pMC1Neo PolyA was first modified to destroy the NcoI site located in the Neo<sup>R</sup> gene and to remove the BamHI, HincII and SalI sites located at the 3' end of the Neo<sup>R</sup> gene. The EF1 $\alpha$  expression cassette was cloned as an HindIII fragment (end

- 29 -

repaired) into the XhoI (end repaired) site in the modified pMC1Neo PolyA plasmid. A vector with the EF1 $\alpha$  promoter and the TK promoter (which drives expression of the Neo<sup>R</sup> gene) facing in opposite directions was chosen for the expression of CPG2\*; the vector is referred to as pMCEF-Plink. CPG2\* was cloned from pMLVCPG2\* as an NcoI/XbaI fragment into the NcoI/XbaI sites of pMCEF-Plink; this plasmid is referred to as pMCEFCPG2\*. As a control plasmid, the lacZ gene was also cloned into the NcoI/XbaI sites pMCEF-Plink as a NcoI/XbaI fragment from the plasmid MLVBlacZ; this is referred to as pMCEFlacZ.

## 2. Preparation of stable cell lines

Stable cell lines were prepared with NIH3T3 cells to constitutively express CPG2\*, or as a control the lacZ gene product ( $\beta$ -galactosidase). In order to do this, NIH3T3 cells were transfected with pMCEF-CPG2\* or pMCEFlacZ as described in Preparative Example 1, and two days after transfection, the cells were plated at low density into medium containing G418 at 1mg/ml. Colonies coming from single cells could be observed about two weeks later and these were cloned and grown up individually. The expression of CPG2\* was determined by immunoprotein blotting of 30 $\mu$ l of cell extract (extracted as described in Preparative Example 1) from each of the G418 resistant colonies with a CPG2 specific polyclonal antibody. The cell line #9 which expresses high levels of CPG2\* (Figure 3A) and cell line #3, which expresses the  $\beta$ -galactosidase (see below) were selected for further investigation. The expression of CPG2\* and  $\beta$ -galactosidase were verified in the cell lines, by enzymatic assay. For these assays, 1x10<sup>5</sup> cells for lines #3 and #9 were plated into 35mm tissue culture dishes and incubated for 4 days. Cell extracts were prepared as described in Preparative Example 1 and 5 $\mu$ g of protein was subjected to a CMDA degradation assay and a  $\beta$ -galactosidase assay (the CMDA assay is described in Preparative Example 1). For the assay of  $\beta$ -galactosidase activity in cells, 5 $\mu$ g of extracted protein was added to 600  $\mu$ l of assay buffer (40mM Na<sub>2</sub>PO<sub>4</sub>, 26.7mM NaH<sub>2</sub>PO<sub>4</sub>, 6.7mMKCL, 1mMMgSO<sub>4</sub>, 25mM 2-mercaptoethanol, 50mM Tris.HCL, 0.06%v/v Triton X-100, 2.2mM o-nitrophenyl  $\beta$ -D-galactopyranoside) and incubated at 37°C for 60 min. 250 $\mu$ l of 1MNa<sub>2</sub>CO<sub>3</sub> was added to the sample and the OD<sub>420</sub> was measured. The results in Figures 3B and 3C show that extracts from line #3 (open

- 30 -

bars) contain high levels of  $\beta$ -galactosidase activity but no CPG2 activity. The extract from line #9 does not contain any detectable  $\beta$ -galactosidase activity but does contain detectable levels of CPG2 activity.

#### EXAMPLE 2

Cell lines #3 and #9 obtained in Preparative Example 2 were tested for cytotoxicity with the CMDA pro-drug.  $1 \times 10^5$  cells were plated into each well of 24 well dishes, incubated for 40 hours and then treated with increasing concentrations of CMDA at 37°C for 60 minutes. After a further 6 hour incubation, 10% of the cells were re-plated into 24 well dishes, incubated for a further 5 days and cell survival was determined by [ $^3\text{H}$ ]-thymidine incorporation. The results in Figure 4 show that cell line #3 (filled symbols) is not sensitive to even the highest concentration of CMDA tested, whereas #9 (open symbols) is sensitive in a dose dependent manner, showing an IC50 of - 125 $\mu\text{M}$ .

#### EXAMPLE 3

The bystander effect was determined by mixing cell lines #3 and #9 obtained in Preparative Example 2 at a variety of concentrations as indicated and a total of  $1 \times 10^5$  cells were plated per well in 24 well plates and incubated overnight. The cells were treated with 250 $\mu\text{M}$  CMDA for 60 minutes as indicated and cell survival was determined as in Example 2. The results in Figure 5 show that with these conditions, more than 95% of the cells are killed when only 5% of the total cell population express CPG2\*: thus demonstrating that there is a substantial bystander effect with this system.



- 31 -

## SEQUENCE INFORMATION

SEQ ID NO. 1: Genomic DNA and protein sequence of CPG2

ATCATGGATC CAAGCACTGA AGGOGGOGGG CAAGACGGGC GGOGTGGOGA CGCTGTGCAT	60
CGGOGGGGGC GAAGGCAOOG CAGTGGCACT OGAATTGCTA TAAGAAOCAT GGCTGGGGAC	120
GOOOGACAAC AGGOGTOCAC CAGCTTTTTT CATTOOGACA ACCOGAOCGA ACAATGGGTA	180
GAGCAGGAGA TTCC ATG CGC CCA TOC ATC CAC CGC ACA GGC ATC GGC GGC	230
Met Arg Pro Ser Ile His Arg Thr Ala Ile Ala Ala	
1 5 10	
GTG CTG GGC AOC GGC TTC GTG GOG GGC AOC GGC CTG GGC CAG AAG CGC	278
Val Leu Ala Thr Ala Phe Val Ala Gly Thr Ala Leu Ala Gln Lys Arg	
15 20 25	
GAC AAC GTG CTG TTC CAG GCA GCT AOC GAC GAG CAG OCG GGC GTG ATC	326
Asp Asn Val Leu Phe Gln Ala Ala Thr Asp Glu Gln Pro Ala Val Ile	
30 35 40	
AAG ACG CTG GAG AAG CTG GTC AAC ATC GAG ACC GGC ACC GGT GAC GGC	374
Lys Thr Leu Glu Lys Leu Val Asn Ile Glu Thr Gly Thr Gly Asp Ala	
45 50 55 60	
GAG GGC ATC GGC GCT GOG GGC AAC TTC CTC GAG GGC GAG CTC AAG AAC	422
Glu Gly Ile Ala Ala Ala Gly Asn Phe Leu Glu Ala Glu Leu Lys Asn	
65 70 75	
CTC GGC TTC ACG GTC ACG CGA AGC AAG TOG GGC GGC CTG GTG GTG GGC	470
Leu Gly Phe Thr Val Thr Arg Ser Lys Ser Ala Gly Leu Val Val Gly	
80 85 90	
GAC AAC ATC GTG GGC AAG ATC AAG GGC CGC GGC GGC AAG AAC CTG CTG	518
Asp Asn Ile Val Gly Lys Ile Lys Gly Arg Gly Gly Lys Asn Leu Leu	
95 100 105	
CTG ATG TOG CAC ATG GAC AOC GTC TAC CTC AAG GGC ATT CTC GCG AAG	566
Leu Met Ser His Met Asp Thr Val Tyr Leu Lys Gly Ile Leu Ala Lys	
110 115 120	
GCC OCG TTC CGC GTC GAA GGC GAC AAG GGC TAC GGC OCG GGC ATC GGC	614
Ala Pro Phe Arg Val Glu Gly Asp Lys Ala Tyr Gly Pro Gly Ile Ala	
125 130 135 140	
GAC GAC AAG GGC GGC AAC GCG GTC ATC CTG CAC ACG CTC AAG CTG CTG	662
Asp Asp Lys Gly Gly Asn Ala Val Ile Leu His Thr Leu Lys Leu Leu	
145 150 155	
AAG GAA TAC GGC GTG CGC GAC TAC GGC ACC ATC ACC GTG CTG TTC AAC	710
Lys Glu Tyr Gly Val Arg Asp Tyr Gly Thr Ile Thr Val Leu Phe Asn	
160 165 170	
ACC GAC GAG GAA AAG GGT TOC TTC GGC TOG CGC GAC CTG ATC CAG GAA	758
Thr Asp Glu Glu Lys Gly Ser Phe Gly Ser Arg Asp Leu Ile Gln Glu	

- 32 -

175	180	185	
GAA GGC AAG CTG GGC GAC TAC GTG CTC TOC TTC GAG CCC ACC AGC GCA Glu Ala Lys Leu Ala Asp Tyr Val Leu Ser Phe Glu Pro Thr Ser Ala 190 195 200			806
GGC GAC GAA AAA CTC TOG CTG GGC ACC TOG GGC ATC GCC TAC GTG CAG Gly Asp Glu Lys Leu Ser Leu Gly Thr Ser Gly Ile Ala Tyr Val Gln 205 210 215 220			854
GTC AAC ATC ACC GGC AAG GGC TOG CAT GGC GGC GGC GGC GAG CTG Val Asn Ile Thr Gly Lys Ala Ser His Ala Gly Ala Ala Pro Glu Leu 225 230 235			902
GGC GTG AAC GGC CTG GTC GAG GCT TOC GAC CTC GTG CTG GGC ACC ATG Gly Val Asn Ala Leu Val Glu Ala Ser Asp Leu Val Leu Arg Thr Met 240 245 250			950
AAC ATC GAC GAC AAG GGC AAG AAC CTG GGC TTC AAC TGG ACC ATC GGC Asn Ile Asp Asp Lys Ala Lys Asn Leu Arg Phe Asn Trp Thr Ile Ala 255 260 265			998
AAG GGC GGC AAC GTC TOG AAC ATC ATC CCC GGC AGC GGC ACC CTG AAC Lys Ala Gly Asn Val Ser Asn Ile Ile Pro Ala Ser Ala Thr Leu Asn 270 275 280			1046
GGC GAC GTG GGC TAC GGC GGC AAC GAG GAC TTC GAC GGC GGC ATG AAG Ala Asp Val Arg Tyr Ala Arg Asn Glu Asp Phe Asp Ala Ala Met Lys 285 290 295 300			1094
ACG CTG GAA GAG GGC GGC CAG CAG AAG AAG CTG CCC GAG GGC GAC GTG Thr Leu Glu Glu Arg Ala Gln Gln Lys Lys Leu Pro Glu Ala Asp Val 305 310 315			1142
AAG GTG ATC GTC ACG GGC GGC GGC GGC TTC AAT GGC GGC GAA GGC Lys Val Ile Val Thr Arg Gly Arg Pro Ala Phe Asn Ala Gly Glu Gly 320 325 330			1190
GGC AAG AAG CTG GTC GAC AAG GGC GTG GGC TAC TAC AAG GAA GGC GGC Gly Lys Lys Leu Val Asp Lys Ala Val Ala Tyr Tyr Lys Glu Ala Gly 335 340 345			1238
GGC ACG CTG GGC GTG GAA GAG GGC ACC GGC GGC GGC ACC GAC GGC GGC Gly Thr Leu Gly Val Glu Glu Arg Thr Gly Gly Thr Asp Ala Ala 350 355 360			1286
TAC GGC GGC CTC TCA GGC AAG CCA GTG ATC GAG AGC CTG GGC CTG GGC Tyr Ala Ala Leu Ser Gly Lys Pro Val Ile Glu Ser Leu Gly Leu Pro 365 370 375 380			1334
GGC TTC GGC TAC CAC AGC GAC AAG GGC GAG TAC GTG GAC ATC AGC GGC Gly Phe Gly Tyr His Ser Asp Lys Ala Glu Tyr Val Asp Ile Ser Ala 385 390 395			1382
ATT CCG GGC GGC CTG TAC ATG GCT GGC GGC CTG ATC ATG GAT CTG GGC Ile Pro Arg Arg Leu Tyr Met Ala Ala Arg Leu Ile Met Asp Leu Gly 400 405 410			1430

- 33 -

GOC GGC AAG T GAATGCTGOC OOOGGCTTT TCACTOGOGT TGCTOGTGA 1480  
Ala Gly Lys  
415

ACTOCACCCOOC COGAGGGGGA GGOGGGTTC GOCITGGGGC GGGOOGGGGG OGACGGCTC 1540

GTACACATAGA AGGAAGTGC ATGTGTGTA CAGCAGACCA GGAAGOCATC OGOGAOGGG 1600

TGOGGACIT CTGCAAGOC GAACTCTGGC CCAAGGOGC GAATGGGGAC OGOGAGCACA 1660

GCTTTCCCA GAGGCCACA GGOGTGGC TGGGTAGC AGTCTGGTG OOGAGGAGC 1720

ATGGGGGGC OGGCTOGAC TACCTCACT OGGCTGGTG CTGGAGGAGA TGGGGGGG 1780

OGAOGGGGC ACCAGCAOG CCATCAGGT GACCAACTGC CCGTCAAG CCATCTCAT 1840

GCGTACGGC AAGGGCAGC AGAAGAAGCA GTGGCTGAG OGGCTGGGC AGGGGGGAT 1900

GCTGGGGOC TTCTGCTGA CCGAGGCGA GGOGGCAGC GATGCATGA GCTGGGCAC 1960

CAOGGGGGC AAGGAOGGG ACGGCTAGT GATGACGGC GTGAAGCAGT TCATCAACAG 2020

OGGCAAGAAC GGOCAGGTG OGGGATC 2048

- 34 -

CLAIMS

1. A two component system for gene-directed enzyme prodrug therapy which comprises: (a) a vector which encodes a carboxypeptidase which is expressed within a cell; and (b) a prodrug which can be converted into an active drug by said carboxypeptidase.
2. A system according to claim 1 wherein the enzyme is a bacterial carboxypeptidase.
3. A system according to claim 2 wherein the carboxypeptidase is carboxypeptidase CPG2.
4. A system according to any one of claims 1 to 3 wherein the prodrug is a nitrogen mustard prodrug.
5. A system according to any one of claims 1 to 4 wherein the vector comprises a promoter capable of being expressed in a tissue restricted manner.
6. A system according to claim 5 wherein the promoter is a c-erbB2 promoter.
7. A system according to any one of claims 1 to 6 for use in a method of treatment or therapy of the human or animal body.
8. A method of treating a tumour in a patient in need of treatment which comprises administering to said patient an effective amount of a vector encoding an enzyme capable of being expressed on the surface of a cell and a prodrug capable of being converted by said enzyme to an active drug.

1/3

Fig.1A.





Fig.3A.



$2/3$ 

Fig.1B.



CPG2 ENZYME ACTIVITY ASSAY

Fig.2.



3/3

Fig.3.




Fig.4.





Fig.5.



**THIS PAGE BLANK (USPTO)**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**